arXiv:1704.00089v1 [math.RT] 1 Apr 2017

Character Formulas from Matrix Factorisations

Kiran Luecke with an appendix by Constantin Teleman

28 Feb, 2017

Abstract

In the spirit of Freed, Hopkins, and Teleman I establish an equivalence between the category of discrete series representations of a real semisimple Lie group G and a category of equivariant matrix factorisations on a subset of $\mathfrak{g}^*$, in analogy with the situation in [FT] which treated the case when G is compact or a loop group thereof. The equivalence is implemented by a version of the Dirac operator used in [FHT1-3], squaring to the superpotential W defining the matrix factorisations. Using the structure of the resulting matrix factorisation category as developed in [FT] I deduce the Kirillov character formula for compact Lie groups and the Rossman character formula for the discrete series of a real semi-simple Lie group. The proofs are a calculation of Chern characters and use the Dirac family constructed in [FHT1-3]. Indeed, the main theorems of [FHT3] and [FT] are a categorification of the Kirillov correspondence, and this paper establishes that this correspondence can be recovered at the level of characters.

1 Introduction and Main Results

Let V be a representation of a Lie group G. The phrase *Kirillov character formula* is used ubiquitously to refer to any formula which exhibits the character of an irreducible representation of G as a Fourier transform of a coadjoint orbit in $\mathfrak{g}^*$.

Suppose G is compact and connected, and $(V_{-\lambda}, \pi)$ is an irreducible representation of lowest weight $-\lambda$. I give a proof of the well known

Theorem 1: (Kirillov)

$$\mathrm{Tr}_{V_{-\lambda}}(e^X) = \hat{\mathcal{A}}(X) \int_{\mathfrak{g}^*} \delta_{O_{-\lambda-\rho}}(\xi) e^{i\langle \xi, X\rangle} d\xi$$

The proof will be a comparison of Chern characters of super-bundles over subsets of $\mathfrak{g}^*$, calculated explicitly using a 1-parameter family of super-connections whose limits as the parameter tends to 0 and ∞ will be integrated to give the two sides of the Kirillov character formula. The proof is completed by observing the invariance of the Chern character, up to an exact differential form, of a 1-parameter family of super-connections on a bundle. The bundle and super-connections are derived directly from the matrix factorisations constructed by Freed, Hopkins, and Teleman in [FHT1-3] and [FT]. They construct an equivariant family of odd Fredholm operators parametrised by $\mu \in \mathfrak{g}^*$ and show that, when coupled to an irreducible representation $V_{-\lambda}$, the family has kernel only over a single coadjoint orbit $O_{-\lambda-\rho}$ where the kernels assemble to form the natural line bundle $\mathcal{L}(-\lambda-\rho)$ tensored with the Atiyah-Bott-Shapiro-Thom class of the normal bundle of $O_{-\lambda-\rho} \in \mathfrak{g}^*$. Since the K-theory class represented by this Fredholm family is invariant under the compact deformation which scales the orbit to 0, they obtain an Dirac version of the Thom isomorphism $K^0_G(0) \cong K_G^{\dim \mathfrak{g}}(\mathfrak{g}^*)$. Applying

equivariant K-theoretic Chern characters gives the Kirillov formula. In fact, the main theorems in [FHT3] and [FT] should be interpreted as a categorification of the Kirillov correspondence, and this paper I extract that correspondence on the level of functions in a more robust way than the K-theoretic argument above, so that it applies to the non-compact setting below, where such methods break down.

Now I generalise and let G be a connected real semisimple Lie group with maximal compact subgroup K, a Cartan subgroup T, and a Cartan decomposition of the Lie algebra $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. This decomposition is orthogonal for the Killing form[1] $\langle \cdot, \cdot \rangle$. Recall that the *discrete series* representations form a family[2] of unitary representations of G, defined by the fact that all matrix coefficients are in $L^2(G)$. A well-known result of Harish-Chandra states that this family is empty unless rank$G$=rank$K$. Fix an irreducible discrete series representation $(\mathcal{H}_{-\Lambda}, \pi)$. Let $\varphi \in C_c^\infty(U)$ where U is an open neighbourhood of $0 \in \mathfrak{g}$ for which $\exp : U \to G$ is invertible analytic, and define the smeared operator $\pi(\varphi) := \int_{\mathfrak{g}} dX \ \varphi(X) e^X$. This operator is trace class and

Theorem 2: (Rossman)

$$\mathrm{Tr}_{\mathcal{H}_{-\Lambda}}\Big(\int_{\mathfrak{g}} dX \ \varphi(X) e^X \Big) = \int_{\mathfrak{g}^*} \delta_{O_{-\Lambda-\tilde{\rho}}}(\nu) d\xi \int_{\mathfrak{g}} dX \ e^{i\langle \nu, X \rangle} \varphi(X) \hat{\mathcal{A}}(X)$$

To prove this formula I will apply the same Chern character method used above, after extending the required results of [FHT1-3] and [FT] to the discrete series setting: Let $\mathcal{E} \subset \mathfrak{g}$ be the cone of elliptic elements. Denote by $\langle \cdot, \cdot \rangle_0$ the (Riemannian) metric on $\mathcal{E}$ obtained by flipping the sign of the Killing form on $\mathfrak{k}$ in the tangent space $T_X\mathfrak{g} \cong \mathfrak{g}$ to elements $X \in \mathfrak{k}$, and then extending to all of $\mathcal{E}$ by G-equivariance. Denote by $\mathcal{S}_0$ the irreducible $\mathbb{Z}_2$-graded spinor representation of $\mathrm{Cliff}^c(\mathfrak{g}^*, \langle \cdot, \cdot \rangle_0)$. Let $\mathfrak{S}\mathrm{Rep}G_{DS}$ denote the supercategory of $\mathbb{Z}_2$-graded representations of the crossed product algebra $G \ltimes \mathrm{Cliff}^c(\mathfrak{g}^*, \langle \cdot, \cdot \rangle_0)$ such that G acts by a direct sum of discrete series representations. Let $\{e_a\}$ be a basis of $\mathfrak{g}$ with dual basis $\{e^a\}$ and let $W = -ie_a(\delta_1) \otimes e^a + \frac{1}{2} e_a e^a$.

Theorem A: *There is an equivalence of supercategories*

$$\mathfrak{S}\mathrm{Rep}G_{DS} \cong \mathrm{MF}_G(\mathcal{E}, W)$$

Theorem A will follow by combining explicit calculations with certain families of Dirac operators with the results of the Appendix, written by Constantin Teleman.

The outline of the paper is as follows: in Section 1 I prove Theorem 1, the Kirillov character formula for irreducible representations of compact connected Lie Groups via the method of Chern characters outlined above. In Section 2 I construct a family of Dirac operators that will exhibit the equivalence of Theorem A. In Section 3 I combine the method of Chern characters of Section 1 with the results of Section 2 to establish the Rossman character formula for discrete series representations of connected real semisimple Lie groups. The Appendix provides technical results needed to turn the Dirac families of Section 2 into honest matrix factorisations as stated in Theorem A.

2 Section 1: Character Formula for Compact Groups

Throughout this section, G is a compact connected Lie group with an irreducible representation V of lowest weight $-\lambda$.

—

[1] We us the standard convention in which this form is negative definite on the first summand and positive definite on the second.

[2] As the name suggests, this family is indexed by discrete parameters.

Consider the equivariant super-bundle $\mathcal{V} := V \otimes \mathcal{S} \times \mathfrak{g}^* \to \mathfrak{g}^*$, where $\mathcal{S}$ is an irreducible $\mathbb{Z}_2$-graded complex $\mathrm{Cliff}^c(\mathfrak{g}^*)$-module[3]. G acts by the coadjoint representation in the base and the obvious action in the fiber. Let $\gamma(\xi)$ denote Clifford multiplication by $\xi \in \mathfrak{g}^*$, and let $\iota(X)$ denote contraction with the vector field. The trivial connection on $V \times \mathfrak{g}^* \to \mathfrak{g}^*$ gives rise to a super-connection on $\mathcal{V}$, i.e. an odd differential operator on bundle-valued forms

$$A : \Omega^\bullet(\mathfrak{g}^*, \mathcal{V})^\pm \longrightarrow \Omega^\bullet(\mathfrak{g}^*, \mathcal{V})^\mp$$

$$A = d + i\gamma(\xi)$$

and a corresponding *equivariant* super-connection on bundle-valued equivariant forms

$$A : (\mathbb{C}[\mathfrak{g}] \otimes \Omega^\bullet(\mathfrak{g}^*, \mathcal{V})^G)^\pm \longrightarrow (\mathbb{C}[\mathfrak{g}] \otimes \Omega^\bullet(\mathfrak{g}^*, \mathcal{V})^G)^\mp$$

usually written at an element $X \in \mathfrak{g}$ as

$$A(X) := A - \iota(X)$$

Note that the total grading on the two complexes above (with respect to which A_1 is required to be odd) is defined to be the sum of the mod-2 (equivariant) cohomological grading on (equivariant) differential forms and the super-grading on $\mathcal{V}^\pm$. Let $\vec{\mu}(X) := \pounds(X) - [\iota(X), A]$ be the moment of X, given by the difference between the Lie derivative on the total space and parallel super-transport along the base. The G-equivariant Chern character for a test element $X \in \mathfrak{g}$ is defined following Mathai-Quillen as

$$\mathrm{str}(e^{A(X)^2}) := \mathrm{str}(e^{(A)^2 + \vec{\mu}(X)}) = \mathrm{str}(e^{-|\xi|^2 + id\gamma(\xi) + \vec{\mu}(X)})$$

To calculate the integral of this differential form, I use the following

Deformation Lemma: ([BVG] Theorem 7.7) Under 1-parameter deformations of the super-connection, the equivariant Chern character differs by an equivariantly exact differential form.

Proof: Indeed, suppose $A_\epsilon(X)$ is a 1-parameter family of equivariant superconnections. Then

$$\frac{d}{d\epsilon}\mathrm{str}(e^{A_\epsilon(X)^2}) = \mathrm{str}\big([A_\epsilon(X), \frac{dA_\epsilon}{d\epsilon}(X)]e^{A_\epsilon(X)^2}\big)$$

Using the equivariant Bianchi identity $[A_\epsilon(X), A_\epsilon(X)^2] = 0$ this becomes

$$= \mathrm{str}\big([A_\epsilon(X), \frac{dA_\epsilon}{d\epsilon}(X)e^{A_\epsilon(X)^2}]\big)$$

Let d_G denote the differential on equivariant forms. Then we have

$$= d_G\Big(\mathrm{str}(\frac{dA_\epsilon}{d\epsilon}(X)e^{A_\epsilon(X)^2})\Big) \quad \square$$

Consider the family of equivariant superconnections

$$A_\epsilon(X) := d + i\sqrt{\epsilon}\gamma(\xi) - \iota(X)$$

[3]See [FHT2 Prop 1.18]

Then $A_{\epsilon=1}$ coincides with A from before, and

$$\mathrm{str}(e^{A_\epsilon(X)^2}) = \mathrm{str}(e^{-\epsilon|\xi|^2 + i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(X)})$$

Applying the deformation lemma, Stokes theorem, and observing the Gaussian decay of the integrand gives, for any ϵ_1, ϵ_2,

$$\int_{\mathfrak{g}^*} \mathrm{str}(e^{A_{\epsilon_1}(X)^2}) - \mathrm{str}(e^{A_{\epsilon_2}(X)^2}) = \int_{S^{n-1}(r\to\infty)} \int_{\epsilon_1}^{\epsilon_2} \mathrm{str}(\frac{i}{2\sqrt{\epsilon}}\gamma(\xi)e^{A_\epsilon(X)^2})d\epsilon = 0$$

Therefore, to calculate $\int_{\mathfrak{g}^*} \mathrm{str}(e^{A_\epsilon(X)^2})$ it suffices to understand things as $\epsilon \to \infty$, where the Gaussian decay localises things to $\xi = 0$.

Following the notation in [FHT1-3], let $R(X)$ and $\sigma(X)$ denote the Lie algebra action on V and $\mathcal{S}$, respectively. In a basis $\{e^j\}$ of $\mathfrak{g}^*$ with dual basis $\{e^j\}$ one has, at $\xi = 0$

$$i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(X) = i\sum_{j=1}^{n} d\xi_j \gamma(e^j) + X^j(R(e_j) + \sigma(e_j))$$

Calculating the degree $n := \dim G$ part of $\mathrm{str}(e^{A_\epsilon(X)^2})$ by expanding the exponential and taking supertraces gives[4]

$$\lim_{\epsilon\to\infty} \mathrm{str}(e^{-\epsilon|\xi|^2 + i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(X)}) = \delta_0(\xi)d\xi\hat{\mathcal{A}}^{-1}(X)\mathrm{Tr}_V(e^X)$$

which means that

$$\int_{\mathfrak{g}^*} \mathrm{str}(e^{A_\epsilon(X)^2}) = \hat{\mathcal{A}}^{-1}(X)\mathrm{Tr}_V(e^X)$$

–

Now I define a new family of superconnections, which deforms the previous one, and with respect to which the Chern character will give the orbital integral on the right side of the Kirillov formula.

Let $\not{D}_0$ be Kostant's cubic Dirac operator as in [FHT1-3], an endomorphism of $V \otimes \mathcal{S}$ given by

$$\not{D}_0 = \big(R(e_j) + \frac{1}{3}\sigma(e_j)\big) \otimes \gamma(e^j)$$

Consider the 2-parameter family of equivariant super-connections

$$A^t_\epsilon(X) := d + \sqrt{t}\not{D}_0 + i\sqrt{\epsilon}\gamma(\xi) - \iota(X)$$

Then, letting $T(\xi)$ be defined by $T(e^j) := R(e_j) + \sigma(e_j)$ for all j,

$$\mathrm{str}(e^{A^t_\epsilon(X)^2}) = \mathrm{str}(e^{t\not{D}_0^2 + 2i\sqrt{t\epsilon}T(\xi) - \epsilon|\xi|^2 + i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(X)})$$

At $t = 0$ we recover the previous family. Let $\epsilon = t$. As before, the deformation lemma and the Gaussian decay of the resulting coboundary gives invariance of the Chern character, which allows us to consider things when $\epsilon = t \to \infty$. Then the super-trace localises to the kernel of $(\not{D}_0 + i\gamma(\xi))^2$, which is calculated in [FHT3] to be nontrivial only when ξ is in the orbit $O_{-\lambda-\rho}$, and cuts out a

[4] see [BGV] Chapter 7.7 for a sample calculation in low dimensions. Morally, the $\hat{\mathcal{A}}$-genus appears when taking supertraces in the spin representation because of the difference between exponentiating in an exterior algebra and a Clifford algebra.

sub-bundle isomorphic to $\mathcal{L}(-\lambda-\rho)\otimes\mathcal{S}_N$, the natural line bundle tensored with the spinors on the normal bundle in $\mathfrak{g}^*$. Let $\Lambda(V\otimes\mathcal{S})$ be the graded set of weights (with multiplicities) of the graded representation $V\otimes\mathcal{S}$ and $|\tau_\pm\rangle$, $\tau_\pm\in\Lambda(V\otimes\mathcal{S})$ a corresponding weight basis[5]. Then in a transverse slice to the orbit at ξ given by the Cartan subalgebra stabilising it, the Chern character becomes

$$= e^{-|\sqrt{t}(\lambda+\rho)+\sqrt{\epsilon}\xi|^2}\epsilon^{r/2}d\xi^1\wedge ...\wedge\ d\xi^d\sum_{\tau_\pm\in\Lambda(V\otimes\mathcal{S})}\pm e^{2i\sqrt{t\epsilon}\langle\xi,\tau_\pm+\lambda+\rho\rangle}\langle\tau_\pm|e^{\vec{\mu}(X)}|\tau_\pm\rangle$$

The coordinates $d\xi^1,...,d\xi^r$ are chosen to span the Cartan subalgebra, which is the only term in the expanded exponential of $id\gamma(\xi)$ that survives the supertrace over normal spinors $\mathcal{S}_N$. Therefore

$$\lim_{t=\epsilon\to\infty}\mathrm{str}(e^{A_\epsilon^t(X)^2}) = \delta_{-\lambda-\rho}(\xi)d\xi\langle-\lambda-\rho|e^{\vec{\mu}(X)}|-\lambda-\rho\rangle$$

The bundle formed by $|-\lambda-\rho\rangle$ is exactly $\mathcal{L}(-\lambda-\rho)$, on which $\vec{\mu}(X)$ is well-known to be $e^{i\langle X,\xi\rangle}$. Note that the moment on spinors is trivial. Indeed, since the normal bundle to $O_{-\lambda-\rho}$ at a point is exactly its stabiliser, $\mathcal{S}_N$ is equivariantly trivial[6].

$$\lim_{t=\epsilon\to\infty}\int_{\mathfrak{g}^*}\mathrm{str}(e^{A_\epsilon^t(X)^2}) = \int_{\mathfrak{g}^*}\delta_{O_{-\lambda-\rho}}(\xi)e^{i\langle\xi,X\rangle}d\xi$$

To conclude the proof of the Kirillov formula, choose a smooth path in the (ϵ,t)-plane which starts at $(1,0)$ and becomes $\epsilon=t$ at large values and apply the deformation lemma to conclude:

$$\hat{\mathcal{A}}^{-1}(X)\mathrm{Tr}_V(e^X) = \int_{\mathfrak{g}^*}\delta_{O_{-\lambda-\rho}}(\xi)e^{i\langle\xi,X\rangle}d\xi$$

When G is compact, as was assumed in this section, this equality can be derived directly from (twisted equivariant) K-theoretic arguments (the twisted Thom isomorphism (4.10) in [FHTIII]), with the caveat that the $\hat{\mathcal{A}}$-genus appears on the other side, in accordance with the 'Riemann-Roch' discrepancy between Thom classes in K-theory and cohomology. However, the above analytic proof using the explicit formula for the Dirac operator $\not{D}_\xi$ is more robust because it extends, with a few minor adjustments, to the case when G is non-compact and K-theory is not available.

3 Section 2: Dirac Family for Discrete Series

Now G is a connected real semisimple Lie group. Recall that we have fixed a Cartan subgroup $T\subset K\subset G$ with Lie algebra $\mathfrak{h}$, and assume that T is also a Cartan subgroup of G so that the discrete series is non-empty. Throughout this section $\mu\in\mathfrak{h}^*\subset\mathfrak{k}^*$ will be a regular elliptic element. We choose the system of positive roots $\Delta^+=\Delta_c^+\cup\Delta_n^+$ for which μ is antidominant. Let $\tilde{\rho}=\rho_c-\rho_n$. Let $\mathcal{H}_\Lambda$ be the discrete series rep with Blattner parameter (lowest K-type) $-\Lambda$ and let $\mathcal{S}_0$ be the irreducible $\mathbb{Z}_2$-graded spinor representation of $\mathrm{Cliff}^c(\mathfrak{g}^*,\langle\cdot,\cdot\rangle_0)$. Let $L_{-\Lambda}$ be the lowest weight line of $K_{-\Lambda}$, the (unique) K-representation in $\mathcal{H}_\Lambda$ of this type, and $S_{-\tilde{\rho}}$ the lowest weight space of $\mathfrak{S}_0$. For a basis $\{e^\nu\}$ of $\mathfrak{g}^*$ with dual basis $\{e_\nu\}$ of $\mathfrak{g}$ let $\gamma^v=\gamma(e^\nu)$ denote Clifford multiplication by e^ν, σ_ν the action of e_ν on $\mathcal{S}_0$ by pullback via the coadjoint representation, and let R_ν denote the action

[5]Technically there should be a subscript on $\tau_\pm$ which keeps track of multiplicities; it has been dropped to streamline notation.

[6]This is also the reason that the $\hat{\mathcal{A}}$-genus didn't show up this time. Last time we were localised at the origin, and the normal spinor bundle of the origin is not equivariantly trivial as a representation.

of e_ν on $\mathcal{H}_\Lambda$ (or rather a dense subspace of K-finite vectors). In analogy with [FHTII], define the coupled Dirac operator (see [FHTII]) : for $\mu = \mu_\nu e^\nu \in \mathcal{E}^*$

$$\not{D}_\mu := \gamma^\nu (iR_\nu + \frac{i}{3}\sigma_\nu + \mu_\nu)$$

Proposition 1: *The operator $\not{D}_\mu^2 \in End(\mathcal{H}_\Lambda \otimes \mathcal{S}_0)$ is invertible for all $\mu \in \mathcal{E}^*$ except on the coadjoint orbit $\mathcal{O}_{-\Lambda-\tilde{\rho}}$. For μ in this orbit, $ker\not{D}_\mu = ker\not{D}_\mu^2 = L_{-\Lambda} \otimes S_{-\tilde{\rho}}$. Away from $\mathcal{O}_{-\Lambda-\tilde{\rho}}$ the spectrum of $\not{D}_\mu^2$ is bounded a finite distance away from 0. There is an associated family of bounded Fredholm operators $\not{D}_\mu(1-\not{D}_\mu^2)^{-1/2}$ acting on the Hilbert space $\mathcal{H}_\Lambda \otimes \mathcal{S}_0$ which has all the spectral properties stated for $\not{D}_\mu^2$. This family represents a G-equivariant matrix factorisation over $\mathcal{E}^*$.*

After applying the results of the Appendix we find that the map

$$\mathcal{H}_\Lambda \otimes \mathcal{S}_0 \mapsto \not{D}_\mu(1-\not{D}_\mu^2)^{-1/2}$$

exhibits the equivalence of categories in Theorem A.

–

A Formula for the Dirac Operator

To prove Proposition 1, we give a formula for the diagonalisation of $\not{D}_\mu^2$ along a K-finite basis of $\mathcal{H}_\Lambda \otimes \mathcal{S}_0$. To motivate aforementioned sign-flip in the metric used to define the Clifford algebra, we will begin the proof using Killing form $\langle \cdot, \cdot \rangle$, identify where things go wrong, and make the necessary adjustment to obtain $\langle \cdot, \cdot \rangle_0$. We will need the following

Lemma: *In a discrete series representation $\mathcal{H}_\Lambda$ the negative non-compact roots elements kill $L_{-\Lambda}$, the lowest weight line of the lowest K type Let $v \in L_{-\Lambda}$.*

Proof: Indeed, take $X_{-\beta} \in \mathfrak{g}_{-\beta}$ a root element for some $\beta \in \Delta_n^+$.Then $X_{-\beta} \cdot v = 0$ or $X_{-\beta} \cdot v \in L_{-\Lambda-\beta}$. In the latter case, by applying some (possibly empty) combination of compact roots Δ_c we may conclude that $X_{-\beta} \cdot v$ is contained in a K type of highest weight

$$-\Lambda - \beta + \sum_{\alpha \in \Delta_c^+} n_\alpha \alpha; (n_\alpha \geqslant 0)$$

But by Theorem 9.20 Knapp (restated in terms of lowest weights, using the longest element of W_K, the Weyl group of K) the only possible K-types in such a setting are of the form

$$-\Lambda^{'} = -\Lambda + \sum_{\nu \in \Delta^+ \cup \Delta_c^-} n_\nu \nu; (n_\nu \geqslant 0)$$

Hence it must be that $X_{-\beta} \cdot v = 0$, as desired, because the above formulas disagree on the sign of the non-compact simple root, which appears an odd number of times in β, and an even number of times in any compact root. □

The Spin Module

Let $\mathfrak{n} = \bigoplus_{\alpha\in\Delta^+} \mathfrak{g}^{\mathbb{C}}_\alpha$. Consider the spin module

$$\mathcal{S} \cong \mathcal{S}_{\mathfrak{h}*} \otimes \bigwedge \mathfrak{n}_c \otimes \bigwedge \overline{\mathfrak{n}}_n \otimes \det(\mathfrak{n}_c^*)^{1/2} \otimes \det(\mathfrak{n}_n)^{1/2}$$

Recall two standard facts about Clifford algebras: firstly, an orthogonal decomposition of an inner product space $V = V_1 \oplus V_2$ leads to a decomposition $C(V) \cong C(V_1) \otimes C(V_2)$; secondly, the dual of $C(\mathbb{R})$ with the standard inner product on $\mathbb{R}$ is isomorphic to $C(\mathbb{R})$ with the negative definite one. Thus for $\xi \in \mathfrak{s}^*$, $\gamma(\xi)$ acts contragrediently compared to $\gamma(\mathfrak{k}^*)$. Hence, the spin module for a non-compact Lie algebra will be a direct sum of copies of a representation of lowest (resp. highest) weight $-\tilde{\rho}$ (resp. $\tilde{\rho}$) where $-\tilde{\rho} = \rho_c - \rho_n$. Here we see why $\rho_n - \rho_c$ appears in the Blattner parameter above. Note that Clifford multiplication by the root basis vector e_ν (for ν an positive root index), denoted γ^ν, kills the lowest weight space $S_{-\tilde{\rho}}$, which is the lowest weight line in $\bigwedge \mathfrak{n}_c \otimes \bigwedge \overline{\mathfrak{n}}_n \otimes \det(\mathfrak{n}_c^*)^{1/2} \otimes \det(\mathfrak{n}_n)^{1/2}$ tensored with a multiplicity space isomorphic to $\mathcal{S}_{\mathfrak{h}*}$.

Consider the representation $\mathcal{H}_\Lambda \otimes \mathcal{S}$. In the notation of the introduction, we find that on $L_{-\Lambda} \otimes S_{-\tilde{\rho}}$

$$\gamma^{\alpha_c} = R_{-\alpha_c} = \sigma_{-\alpha_c} = 0$$

for $\alpha_c \in \Delta_c$ and

$$\gamma^{\alpha_n} = R_{-\alpha_n} = \sigma_{-\alpha_n} = 0$$

for $\alpha_n \in \Delta_n$.

Using the definition of the Dirac operator given above one finds

$$\not{D}_\mu\big|_{L_{-\Lambda}\otimes S_{-\tilde{\rho}}} = \gamma(\Lambda + \tilde{\rho} + \mu)$$

and so

$$\not{D}_\mu^2\Big|_{L_{-\Lambda}\otimes S_{-\tilde{\rho}}} = -|\Lambda + \tilde{\rho} + \mu|^2$$

meaning that $\not{D}_\mu$ has kernel on this weight space only when $\mu = -\Lambda - \tilde{\rho}$.

To rule out kernel on other weight spaces one would like to mimic the argument given in [FHTII] for the compact case - namely, use the commutation relations of $\not{D}_\mu$ to prove that $\not{D}_\mu^2$ is a nonpositive operator with maximum on $L_{-\Lambda} \otimes S_{-\tilde{\rho}}$. This is done as follows. For $X \in \mathfrak{g}$ define the μ-shifted Lie algebra action $\pi(X) = X^a(R_a + \sigma_a - i\mu_a)$. One calculates[7]

$$[\not{D}_\mu, \pi(\xi_*)] = \gamma\Big([\mu_*, \xi_*]^*\Big)$$

$$[\not{D}_\mu, \gamma(\xi)] = -2i\pi(\xi_*)$$

Here $\eta \mapsto \eta_*$ is the map $\mathfrak{g}^* \to \mathfrak{g}$ induced by the Killing form, and $\eta \mapsto \eta^*$ is its inverse.

Iterating the commutators gives, $\forall \xi \in \mathfrak{g}^*$:

$$[\not{D}_\mu^2, \pi(\xi_*)] = [2T(\xi_*) - |\mu|^2, \pi(\xi_*)]$$

$$[\not{D}_\mu^2, \gamma(\xi)] = [2T(\xi_*) - |\mu|^2, \gamma(\xi)]$$

Hence, for $\mu = 0$, one has

$$[\not{D}_0^2, \pi(\xi_*)] = [\not{D}_0^2, \gamma(\xi)] = 0$$

[7] These calculations are identical to the compact case, see [FHTII] Prop. 1.19

Therefore, $\not{D}_0^2$ is in the center of the algebra $G \rtimes \mathrm{Cliff}(\mathfrak{g}^*)$. These operators generate $\mathcal{H} \otimes \mathcal{S}$ from $L_{-\Lambda-\tilde{\rho}}$, so $\not{D}_0^2$ a constant. This constant is easily calculated to be $-|\Lambda+\tilde{\rho}|^2$. But then

$$\not{D}_\mu^2 = \not{D}_0^2 + 2iT(\mu) - |\mu|^2 = -|\Lambda+\tilde{\rho}|^2 + 2iT(\mu) - |\mu|^2$$

where

$$T(\mu) := \mu_*^a(R_a + \sigma_a)$$

When G is compact, so that the representation $V \otimes \mathcal{S}$ is finite dimensional with lowest $T(\mu)$-eigenspace $L_{-\Lambda} \otimes S_{-\tilde{\rho}}$, this formula simultaneously proves the desired nonpositivity and invertibility of $\not{D}_\mu^2$ for $\mu \notin O_{-\lambda-\rho}$, by applying highest/lowest weight theory.

In the current case we cannot use this argument. The discrete series representations are infinite dimensional and there is in general[8] no minimum eigenspace for μ, only a minimal K-type. If the spectrum of $T(\mu)$ is unbounded, there exists a weight space (in some large enough K-type) where the Dirac operator for, say, $\mu = -\Lambda - \tilde{\rho}$, has a positive eigenvalue. Scaling $\mu \mapsto \epsilon\mu$ down to zero inside $\mathfrak{h}^*$, we see that the eigenvalue on this weight space must eventually become very close to $\not{D}_0^2 = -|\Lambda+\tilde{\rho}|^2$ which is negative. It does so continuously; hence at some point this eigenvalue vanishes and we get kernel away from $O_{-\Lambda-\rho}$.

To fix the problem, we must study a slightly different Dirac operator. Namely, we replace the trivial bundle of Clifford algebras $\mathrm{Cliff}^c(\mathfrak{g}^*, \langle\cdot,\cdot\rangle)$ over $\mathfrak{g}^*$ to the bundle with fibre $\mathrm{Cliff}^c(\mathfrak{g}^*, \langle\cdot,\cdot\rangle_0)$, the Clifford algebra on $T\mathfrak{g}^*$ defined by the metric constructed in the Introduction. The Dirac operator with respect to this Clifford algebra builds an equivariant family only when restricted to the elliptic locus $\mathcal{E}^*$. We now write the formula for this Dirac Laplacian and compare it to the previous one. Note that $\not{D}_0^2 = g^{ij}(\mu)(R_iR_j + \frac{1}{3}\sigma_i\sigma_j)$ where $g^{ij}(\mu)$ is the metric in a basis of $T_\mu\mathcal{E}^*$. Choose this basis to respect the Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Let $\mu \in \mathfrak{k}$. Then the Dirac Laplacian with respect to the Killing metric was

$$\not{D}_\mu^2 := \left(\not{D}_0^2\right)_{\mathfrak{k}} + \left(\not{D}_0^2\right)_{\mathfrak{p}} + 2iT(\mu) - |\mu|^2$$

while the new Dirac Laplacian, defined with the metric $\langle\cdot,\cdot\rangle_0$, is

$$\not{D}_\mu^2 := \left(\not{D}_0^2\right)_{\mathfrak{k}} - \left(\not{D}_0^2\right)_{\mathfrak{p}} + 2iT(\mu) - |\mu|^2$$

$$\not{D}_0^2 = \left(\not{D}_0^2\right)_c + \left(\not{D}_0^2\right)_n$$

Determining the behaviour of $\left(\not{D}_0^2\right)_c - \left(\not{D}_0^2\right)_n$ is simple:

$$\begin{aligned}\left(\not{D}_0^2\right)_c - \left(\not{D}_0^2\right)_n &= -\left(\not{D}_0^2\right)_{\mathfrak{k}} - \left(\not{D}_0^2\right)_{\mathfrak{p}} + 2\left(\not{D}_0^2\right)_c \\ &= |\Lambda+\tilde{\rho}|^2 + 2\left(\not{D}_0^2\right)_c\end{aligned}$$

But $\left(\not{D}_\mu^2\right)_c$ is just the Dirac Laplacian for K (at $\mu \in \mathfrak{k}$) whose behaviour diagonalises along K-types and whose behaviour on each K-type $V_{K,\tau}$ we know to be that of a nonpositive operator with maximum eigenvalue on the lowest weight space $W_{-\tau}$subset $V_{K,\tau}$.

[8] An exception is, e.g. $SL_2(\mathbb{R})$, where $K = T$ so the K-types are 1-dimensional and the discrete series really are lowest/highest weight representations.

Hence, under the new definition of the Dirac operator,

$$\not{D}_\mu^2\Big|_{V_{K,\tau}} = |\Lambda+\tilde{\rho}|^2 - |\tau+\rho_c|^2 + \left(\not{D}_\mu^2\right)_c$$

By Knapp's theorem about K-types from above, adapted to account for the presence of spinors, any K-type in $\mathcal{H}\otimes\mathcal{S}$ has highest weight of the form

$$\tau = \Lambda + \tilde{\rho} + \sum_{\beta\in\Delta_n^+} n_\beta\beta + \sum_{\nu\in\Delta^+} c_\nu\nu$$

where $n_\beta \in \mathbb{Z}_{\geqslant 0}$ and $c_\nu \in \{0,1\}$. From this it follows that

$$|\Lambda+\tilde{\rho}|^2 - |\tau+\rho_c|^2 \leqslant 0$$

for every possible τ, which implies that $\not{D}_\mu^2$ is a non-positive operator whose maximum eigenvalue occurs on the lowest weight space $L_{-\Lambda}\otimes S_{-\tilde{\rho}}$ of the lowest K-type in $\mathcal{H}\otimes\mathcal{S}$, and which vanishes, for elliptic elements $\mu\subset\mathfrak{h}^*$, only when $\mu = -\Lambda-\tilde{\rho}$. As mentioned before, the space $L_{-\Lambda}\otimes S_{-\tilde{\rho}}$ is a 1 -dimensional subspace, up to a multiplicity space isomorphic to $\mathcal{S}_{\mathfrak{h}*}$.

–

Bounding the Spectrum Away from 0

In this section we give a lower bound for the square norm of the eigenvalues of $\not{D}_\mu$ away from the subspace $L_{-\Lambda}\otimes S_{-\tilde{\rho}}$ for $\mu\in\mathcal{E}^*$, thereby exhibiting the rigidity of discrete series representations.

We relax notation and do not distinguish between $\mu\in\mathfrak{g}^*$ and its image in $\mathfrak{g}$ under the isomorphism induced by the inner product $\langle\cdot,\cdot\rangle_0$. Note that square norms like $|\mu|^2$ occurring below are taken with this inner product, *not* the Killing form. Proceed case by case:

Case 1: Elliptic - Covered in the last subsection.

Case 3: Nilpotent Let μ be nilpotent. But then $|\mu|^2 = 0$ and $T(\mu)$ is nilpotent, so the spectrum of $\not{D}_\mu^2$ is bounded below in square norm by $|\Lambda+\tilde{\rho}|^2$.

Case 4: Let $\mu = \mu_e + \mu_n$ be mixed elliptic/nilpotent. After possible conjugation we can assume the two elements commute. Then we have

$$\not{D}_\mu^2 = \not{D}_{\mu_e}^2 - 2iT(\mu_n) + |\mu_e|^2 - |\mu_e+\mu_n|^2$$

The two terms in the last line commute and the second is nilpotent, so the expression is strictly negative unless $\mu_n = 0$ and we reduce to Case 1.

Hence, for all $\mu\in\mathcal{E}^*\backslash O_{-\Lambda-\tilde{\rho}}$, the spectrum of the Dirac operator is contained in the complement of an open ball about $0\in\mathbb{C}$ of finite radius.

4 Section 3: Character Formula for Discrete Series

Now let G be a connected real semisimple Lie group with maximal compact subgroup K, and suppose r =rank$G$=rank$K$, so that G has discrete series. Fix an irreducible discrete series representation $(\mathcal{H}_{-\Lambda},\pi)$. Here $-\lambda$ is the lowest weight line of the lowest K-type, which labels the discrete series in an analogous way to the case finite dimensional representations of a compact group (see [K] 9.20). Because $\mathcal{H}_{-\lambda}$ is infinite dimensional, the traces of operators $\pi(g)$, $g\in G$, may not be defined. There is a Kirillov character formula, due to Rossman, for the character of the smeared operator

$\pi(\varphi) := \int_{\mathfrak{g}} dX\ \varphi(X)e^X$ which turns out to be trace class for any $\varphi \in C_c^\infty(U)$ where U is an open neighbourhood of $0 \in \mathfrak{g}$ for which $\exp : U \to G$ is invertible analytic. The formula is:

$$\mathrm{Tr}_{\mathcal{H}_{-\lambda}}\Big(\int_{\mathfrak{g}} dX\ \varphi(X)e^X\Big) = \int_{\mathfrak{g}^*} \delta_{O_{-\Lambda-\tilde{\rho}}}(\nu)d\xi \int_{\mathfrak{g}} dX\ e^{i\langle \nu, X\rangle}\varphi(X)\hat{\mathcal{A}}(X)$$

–

Let $\mathcal{E} \subset \mathfrak{g}^*$ denote the locus of elliptic elements. Consider the super-bundle $\mathcal{H}_{-\Lambda} \otimes \mathcal{S}_0 \times \mathcal{E} \to \mathcal{E}$, on which the discrete series matrix factorisations live ([LT]). The subscript on the spinor factor denotes the fact that the spin representation $\mathcal{S}_0$ is defined for the Clifford algebra on $T^*\mathcal{E}$ with the (Riemannian) metric obtained at points in $\mathfrak{t}^* \subset \mathfrak{k}^*$ after flipping all signs of the Killing form on $\mathfrak{p}$ to be positive, and extending to $\mathcal{E}$ by equivariance. The extension is possible since this bilinear form is invariant under T, the compact Cartan subgroup which stabilises an elliptic element. However, extending to general elements in $\mathfrak{g}^*$ fails because their stabilisers may not fix this deformed version of the Killing form. The reason for flipping the metric in the first place is explained in [LT]; essentially, having passed to a setting where representations have unbounded weights, the flip is required to make the kernel of $\not{D}_0 + i\gamma(\xi)$ mimic the case of compact G and cut out the natural line bundle over a single coadjoint orbit.

Let $A_\epsilon^t(\varphi)$ and $\vec{\mu}(\varphi)$ be defined by the linear extension of the action of $\mathfrak{g}$ on $\mathcal{H}_{-\lambda} \otimes \mathcal{S}_0 \times \mathfrak{g}^* \to \mathfrak{g}^*$ to an action of $C_c^\infty(U)$. For example, $\vec{\mu}(\varphi) := \int_{\mathfrak{g}} dX\ \varphi(X)\vec{\mu}(X)$.

We consider the same family of superconnections as before. Although the deformation lemma continues to hold, a more careful argument is needed to establish invariance of the integral. This is because of the issues with the metric and the fact that $\partial\mathcal{E}$ includes the nilpotent cone $\mathcal{N}$, where there is no Gaussian decay. For the sake of continuity, I will set up all the necessary integrals at key values of (ϵ, t) and the give the invariance arguments at the end.

So the task at hand is to evaluate, once at $(\epsilon \to \infty), t = 0)$ and once at $(\epsilon \to \infty, t - \epsilon)$,

$$\int_{\mathcal{E}} \mathrm{str}(e^{A_\epsilon^t(\varphi)^2}) = \int_{\mathcal{E}} \mathrm{str}(e^{t\not{D}_0^2 + 2i\sqrt{t\epsilon}T(\xi) - \epsilon|\xi|^2 + i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(\varphi)})$$

So in analogy with the compact case we find immediately that

$$\lim_{\epsilon\to\infty, t=0} \int_{\mathcal{E}} \mathrm{str}\Big(e^{A_\epsilon^0(\varphi)^2}\Big) = \int_{\mathcal{E}} \delta_0(\xi)d\xi^1 \wedge ... \wedge d\xi^n \hat{\mathcal{A}}^{-1}(\varphi)\mathrm{Tr}_{\mathcal{H}_{-\Lambda}}(\varphi)$$

$$= \hat{\mathcal{A}}^{-1}(\varphi)\mathrm{Tr}_{\mathcal{H}_{-\lambda}}(\varphi) := \mathrm{Tr}_{\mathcal{H}_{-\Lambda}}\Big(\int_{\mathfrak{g}} dX\ \varphi(X)\hat{\mathcal{A}}(X)^{-1}e^X\Big)$$

It remains to recalculate the Chern character at $(\epsilon \to \infty, t = \epsilon)$. The required spectral properties of $\not{D}_0$ in this setting are established in [LT]. The results are essentially the same as for compact groups - the operator has kernel over a single regular elliptic orbit, and those kernel subspaces assemble to a bundle which is isomorphic to the natural line bundle $\mathcal{L}(-\Lambda - \tilde{\rho}) \otimes \mathcal{S}_N$. Borrow existing notation from Section 1, and letting $K(\tau_\pm)$ denote the square norm of the K-type in which $|\tau_\pm\rangle$ is contained, we have

$$\mathrm{str}(e^{A_\epsilon^t(\varphi)^2}) = \mathrm{str}(e^{t\not{D}_0^2 + 2i\sqrt{t\epsilon}T(\xi) - \epsilon|\xi|_0^2 + i\sqrt{\epsilon}d\gamma(\xi) + \vec{\mu}(\varphi)})$$

$$= e^{-|\sqrt{t}(\Lambda+\tilde{\rho})+\sqrt{\epsilon}\xi|_0^2}\epsilon^{r/2}d\xi^1 \wedge ... \wedge\ d\xi^d \sum_{\tau_\pm \in \Lambda(\mathcal{H}\otimes\mathcal{S})} \pm e^{2i\sqrt{t\epsilon}\langle \xi, \tau_\pm + \lambda + \rho\rangle + t(|\Lambda+\tilde{\rho}|^2 - K(\tau_\pm))}\langle \tau_\pm | e^{\vec{\mu}(\varphi)} | \tau_\pm \rangle$$

There is exponential damping away from $\mathcal{L}(-\Lambda-\tilde{\rho})\otimes\mathcal{S}_N$, where the moment is $\vec{\mu}(\varphi) := \int_{\mathfrak{g}} dX\ \varphi(X)\vec{\mu}(X) = \int_{\mathfrak{g}} dX\ \varphi(X)e^{i\langle\xi,X\rangle}$. Hence

$$\lim_{t=\epsilon\to\infty} \mathrm{str}(e^{A_\epsilon^t(\varphi)^2}) = \delta_{O_{-\Lambda-\tilde{\rho}}}(\xi)d\xi\langle -\Lambda-\tilde{\rho}|e^{\vec{\mu}(\varphi)}|-\lambda-\rho\rangle$$

and so

$$\lim_{t=\epsilon\to\infty}\int_{\mathcal{E}} \mathrm{str}(e^{A_\epsilon^t(\varphi)^2}) = \int_{\mathcal{E}} \delta_{O_{-\Lambda-\tilde{\rho}}}(\xi)e^{i\langle\xi,\varphi\rangle}d\xi = \int_{\mathcal{E}} \delta_{O_{-\Lambda-\tilde{\rho}}}(\xi)d\xi\int_{\mathfrak{g}} dX\ e^{i\langle\nu,X\rangle}\varphi(X)$$

Provided the integral is invariant under 1 parameter deformations, we can compare the integral at the two limits of this family to conclude

$$\mathrm{Tr}_{\mathcal{H}_{-\lambda}}\Big(\int_{\mathfrak{g}} dX\ \varphi(X)\hat{\mathcal{A}}(X)^{-1}e^X\Big) = \int_{\mathcal{E}} \delta_{O_{-\Lambda-\tilde{\rho}}}(\xi)d\xi\int_{\mathfrak{g}} dX\ e^{i\langle\xi,X\rangle}\varphi(X)$$

Replacing $\varphi(X)$ with $\varphi(X)\hat{\mathcal{A}}(X)$ gives Rossman's Kirillov character formula.

To establish invariance of the integral, I work in segments. Let $P(1)$ denote the straight line $(0,1)\to(0,\epsilon)$, $P(2)$ the straight line $(0,1)\to(1,1)$, and $P(3)$ the straight line $(1,1)\to(\epsilon,\epsilon)$. Let $\mathcal{E}(\infty)$ denote the part of $S^{n-1}(r\to\infty)$ bounding the elliptic locus at ∞, and let $\partial\mathcal{E}$ denote the boundary of $\mathcal{E}$ in $\mathfrak{g}^*$. The deformation lemma and Stokes theorem give

$$P(1): \quad \int_{\mathcal{E}} \mathrm{str}(e^{A_\epsilon(\varphi)^2}) - \mathrm{str}(e^{A_0(\varphi)^2}) = \int_{\mathcal{E}(\infty)\cup\partial\mathcal{E}}\int_0^s \mathrm{str}(\frac{i}{2\sqrt{s}}\gamma(\xi)e^{A_s(X)^2})ds$$

$$P(2): \quad \int_{\mathcal{E}} \mathrm{str}(e^{A_1^1(\varphi)^2}) - \mathrm{str}(e^{A_1^1(\varphi)^2}) = \int_{\mathcal{E}(\infty)\cup\partial\mathcal{E}}\int_0^1 \mathrm{str}(\frac{1}{2\sqrt{s}}\not{D}_0 e^{A_1^s(X)^2})ds$$

$$P(3): \quad \int_{\mathcal{E}} \mathrm{str}(e^{A_\epsilon^\epsilon(\varphi)^2}) - \mathrm{str}(e^{A_1^1(\varphi)^2}) = \int_{\mathcal{E}(\infty)\cup\partial\mathcal{E}}\int_1^s \mathrm{str}(\frac{1}{2\sqrt{s}}(\not{D}_0 + i\gamma(\xi))e^{A_s^s(X)^2})ds$$

In all three cases, the integral over $\mathcal{E}(\infty)$ vanishes due to Gaussian decay. For $P(2)$ and $P(3)$, the integrand decays to 0 at $\partial\mathcal{E}$ as well. Indeed, the Gaussian factor is of the form $e^{-|\Lambda+\xi|^2}$, peaked on the (elliptic) orbit of λ and decaying in the stabiliser directions (which are orthogonal to the orbit in current metric) and with respect to which the nilpotent cone is at ∞.[9] It remains to check the $\partial\mathcal{E}$ integral for $P(1)$

$$\int_{\partial\mathcal{E}}\int_0^s \mathrm{str}(\gamma(\xi)e^{i\sqrt{s}d\gamma(\xi)+\vec{\mu}(\varphi)})\frac{ids}{2\sqrt{s}}$$

Change the basis $\{e^j\}$ so that locally the first $p<n$ span $\partial\mathcal{E}$. It suffices to pick out the terms in the integrand, after expanding the exponential, whose cohomological component is a wedge sum of $d\xi_1,...,d\xi_p$. Let $J=(j_1,...,j_p)$ be a multi-index of length p, and let $\gamma^J := \gamma(e^{j_1})...\gamma(e^{j_p})$. Let $\vec{\mu}_J(\varphi)$ denote the contribution of $\vec{\mu}(\varphi)$ to the term in the expansion of the integrand which has $d\xi_J := d\xi_{j_1}...d\xi_{j_p}$ in it. Let $\mathfrak{s}\in S_p$ be a permutation[10]. Then the integral simplifies to

$$\int_{\partial\mathcal{E}}\int_0^s \frac{ids}{2\sqrt{s}}(i\sqrt{s})^p d\xi_1\wedge...\wedge d\xi_p\sum_{j=1}^p\sum_{\mathfrak{s}\in S_p}\xi_j(-1)^{|\mathfrak{s}|}\mathrm{str}\big(\gamma(e^j)\gamma^{\mathfrak{s}\cdot J}\vec{\mu}_{\mathfrak{s}\cdot J}(\varphi)\big)$$

[9] A good picture to keep in mind is the case of $SL_2(\mathbb{R})$, where the elliptic orbit is half of a 2 sheeted hyperboloid, the stabiliser to a point on this hyperboloid is the line through that point and the origin, which only approaches the nilcone as the point moves to ∞ on the orbit.

[10] The standard symbol σ is already taken...

For $j, k \in \{1, ..., p\}$, $\gamma(e^j)$ appears in $\gamma^{\mathfrak{s}\cdot J}$ for all $\mathfrak{s}$ and $[\gamma(e^j), \gamma(e^k)] = 0$, in particular $\gamma(e^j)^2 = 0$. Indeed, the closure of the coadjoint orbit of any element in $\partial\mathcal{E}$ contains the origin. Thus, up to a sign, one can move the two $\gamma(e^j)$'s next to each other, whence they annihilate. Therefore each term in the integrand is 0 and we are done.

5 Appendix: C^∞ Matrix Factorisations

By Constantin Teleman. To appear in the final version.

6 References

[BGV] N. Berline, E. Getlzer, M. Vergne: Heat Kernels and Dirac Operators *Grundlehren Text Edition*, 2004

[FHT1] D. Freed, M. Hopkins, and C. Teleman: Loop Groups and Twisted K-Theory *Journal of Topology* **4**, 2011

[FHT2] D. Freed, M. Hopkins, and C. Teleman: Loop Groups and Twisted K-Theory *Journal of the American Mathematical Society* **26**, 2013

[FHT3] D. Freed, M. Hopkins, and C. Teleman: Loop Groups and Twisted K-Theory *Annals of Math* **174**, 2011.

[FT] D. Freed and C. Teleman: Dirac Families for Loop Groups as Matrix Factorisations *Comptes Rendus de l'Acadademie des Sciences* Paris, **353**, (2015)

[K] A. Knapp: Representation Theory of Semisimple Groups *Princeton Landmarks in Mathematics*, 1997

[N] Y. Nakagama: Spectral Analysis in Krein Spaces *Publications of the Research Institute for Mathematical Sciences, Kyoto University* **24**, 1988

[O] D. Orlov: Triangulated Categories of Singularities and D-branes in Landau-Ginzburg Models *Proceedings of the Steklov Institute for Mathematics* **3**, 2004